arXiv:1607.04030v2 [math.GT] 6 Jan 2017

THE EXTENDED MAPPING CLASS GROUP CAN BE GENERATED BY TWO TORSIONS

XIAOMING DU

Abstract. Let S_g be the closed oriented surface of genus g and let $\mathrm{Mod}^{\pm}(S_g)$ be the extended mapping class group of S_g. When the genus is at least 5, we prove that $\mathrm{Mod}^{\pm}(S_g)$ can be generated by two torsion elements. One of these generators is an order 2 element, and the other one is an order 4g+2 element.

1. Introduction

Let S_g be the closed oriented surface of genus g. The extended mapping class group $\mathrm{Mod}^{\pm}(S_g)$ is defined by $\mathrm{Homeo}^{\pm}(S_g)/\mathrm{Homeo}_0(S_g)$, the group of homotopy classes of homeomorphisms (including orientation-preserving ones and orientation-reversing ones) of S_g, and the mapping class group $\mathrm{Mod}(S_g)$ is defined by $\mathrm{Homeo}^{+}(S_g)/\mathrm{Homeo}_0(S_g)$, the group of orientation-preserving homotopy classes of homeomorphisms of S_g.

For $\mathrm{Mod}(S_g)$, Dehn and Lickorish found independently Dehn twist generating sets of $\mathrm{Mod}(S_g)$ [3] [11]. Humphries reduced the number of Dehn twist generators to the lowest bound [5]. Wajnryb in [17] found that the minimal number of the generators (not only Dehn twist generators) for the mapping class groups is 2.

The torsion elements have finite order. We are also interest in the torsion generating set. McCarthy and Papadopoulos in [15] proved that $\mathrm{Mod}(S_g)$ can be generated by infinitely many elements of order 2 when $g \geq 3$. Luo showed that $\mathrm{Mod}(S_g)$ is generated by 12g+6 elements of order 2 when $g \geq 3$ [12]. Brendle and Farb reduced the number of the involution generators to 6 [1] and prove $\mathrm{Mod}(S_g)$ can be generated by three torsion elements. Kassabov reduced the number of the involution generators to 4 [8] for $g \geq 7$. Korkmaz in [9] proved $\mathrm{Mod}(S_g)$ can be generated by two torsion elements of

2010 *Mathematics Subject Classification.* 57N05, 57M20, 20F38.

Key words and phrases. mapping class group, generator, torsion.

This research is supported by NSFC (Grant No. 11401219). I wish to thank BICMR for its hospitality and thank Boju Jiang who asked me about the problem of finding lower bounds of the orders and the numbers of torsion generators for the mapping class groups. I also wish to thank Dingsen Yan who teach me how to draw pictures with the metapost package of the latex.

order 4g+2. Monden in [14] proved $\mathrm{Mod}(S_g)$ can be generated by 3 torsion elements of order 3.

For the extended mapping class group $\mathrm{Mod}^{\pm}(S_g)$, Dehn-Nielsen-Baer theorem [4] says that $\mathrm{Mod}^{\pm}(S_g)$ is isomorphic to $\mathrm{Out}(\pi_1(S_g))$, the outer automorphism group of $\pi_1(S_g)$. Ivanov proved that $\mathrm{Mod}^{\pm}(S_g)$ is the automorphism group of the curve complex [6]. Brock and Margalit showed that $\mathrm{Mod}^{\pm}(S_g)$ is the automorphism group of the pants complex and it is also the isometry group of the Teichmüller space under the Weil-Petersson metric ([2],[13]). Korkmaz showed that $\mathrm{Mod}^{\pm}(S_g)$ can be generated by 2 elements, one of which is a Dehn twist [9]. Stukow in [16] proved $\mathrm{Mod}^{\pm}(S_g)$ is generated by 3 elements of order 2.

It is an open problem that whether $\mathrm{Mod}^{\pm}(S_g)$ can be generated by two torsion elements (See [10], Problem 5.3). In this paper, we answer this question affirmatively:

Theorem 1.1. *For $g \geq 5$, the extended mapping class group $Mod^{\pm}(S_g)$ can be generated by 2 torsion elements. One of these generators is an order 2 element and the other one is an order 4g+2 element.*

2. Preliminaries

Notations:

(a) We use the convention of functional notation, namely, elements of the mapping class group are applied right to left, i.e. the composition FG means that G is applied first.

(b) A Dehn twist means a right-hand Dehn twist.

(c) We denote the curves by the letters a, b, c, d (possibly with subscripts) and the Dehn twists about them by the corresponding capital letters A, B, C, D. Notationally we do not distinguish a diffeomorphism/curve and its isotopy class.

We recall the following results (see, for instance, [4]):

Lemma 2.1 (Conjugacy relation)**.** *For any $\varphi \in Mod(S_g)$ and any isotopy classes a, b of simple closed curves in S_g satisfying $\varphi(a) = b$, we have:*

$$B = \varphi\, A\, \varphi^{-1}.$$

Lemma 2.2 (Dehn twists along disjoint curves commute)**.** *Let a, b be two simple closed curves on S_g. If a is disjoint from b, then*

$$AB = BA.$$

Let $c_1, c_2, \ldots, c_t$ be simple closed curves on S_g. If $i(c_j, c_{j+1}) = 1$ for all $1 \leq j \leq k-1$ and $i(c_j, c_k) = 0$ for $|j-k| > 1$, then we call $c_1, c_2, \ldots, c_t$ is a *chain*. Take a regular neighbourhood of their union. The boundary of

this neighbourhood consist of one or two simple closed curves, depending on whether t is even or odd. In the odd case, denote the boundary curve by d. In the even case, denote the boundary curves by d_1 and d_2. We have the following *chainrelation*:

Lemma 2.3 (Chain relation)**.**

$$\begin{array}{lcll}(C_tC_{t-1}\dots C_2C_1)^{2t+2} & = & D & t \text{ even},\\ (C_tC_{t-1}\dots C_2C_1)^{t+1} & = & D_1D_2 & t \text{ odd}.\end{array}$$

Dehn discovered the lantern relation and Johnson rediscovered it ([7]). This is a very useful relation in the theory of maping class groups.



Figure 1

Lemma 2.4 (Lantern relation)**.** *Let a, b, c, d, x, y, z be the curves showed in Figure 2 on a genus zero surface with four boundaries. Then*

$$ABCD = XYZ.$$

In other words, since a, b, c are disjoint from x, y, z, we have

$$D = (XA^{-1})(YB^{-1})(ZC^{-1}).$$

The elements having the form of UV^{-1} for some disjoint simple closed curves u, v play a crucial role in our proof. The proof of the main result relies on Humphries' theorem:

Theorem 2.5 (Humphries)**.** *Let $a_1, a_2, \dots a_{2g}, b$ as the curves in the picture on the left side of Figure 2. Then the mapping class group Mod(S_g) is generated by A_i's and B.*



Figure 2

Lemma 2.6. *Let* $\sigma = A_{2g}A_{2g-1}\ldots A_2A_1$. *Then* σ^{4g+2} *equals identity. Moreover,* $\sigma^{-1}(a_i) = a_{i+1}$ *for* $1 \leq i \leq 2g-1$.

Proof. $a_1, a_2, \ldots, a_{2g}$ is a chain and the boundary curve of their regular neighbourhood is a curve that bound a disk. Hence by the chain relation, $(A_{2g}A_{2g-1}\ldots A_2A_1)^{4g+2} = 1$. $\sigma^{-1}(a_i) = a_{i+1}$ follows from the braid relation.

Another viewpoint is the that we can draw the curves a_i's as the picture on the right side of Figure 2, considering the surface of genus g as the $(4g+2)$-gon whose corresponding opposite sides are glued together. σ is the composition of the Dehn twists along each a_i's. So σ gives a $\frac{\pi}{2g+1}$ clockwise rotation of such $(4g+2)$-gon. Obviously, $\sigma^{-1}(a_i) = a_{i+1}$. □

If we denote $a_k = \sigma^{1-k}(a_1)$ for all integer k, then obviously $\sigma^{-1}(a_i) = a_{i+1}$ still holds and $a_{i+2g+1} = a_i$.

Since σ is an order $4g+2$ element. We denote $\sigma^{-k}(b)$ as b_k and $b_0 = b$. Then, similar to the case of the a_i curves, let $b_{i+4g+2} = b_i$ and $\sigma^{-1}(b_i) = b_{i+1}$. Figure 3 shows the images of b_i's on the surface.



Figure 3

3. The main result and the proof

Let τ be the symmetry of the $(4g+2)$-gon as in Figure 4. Then τ induces an orientation-reversing homeomorphism of the genus g surface. We still denote the homotopy class of such a homeomorphism as τ. Since τ preserves the curve b but reverse the direction of b, it is easy to check that $(\tau \circ B)^2 = Id$.



Figure 4

Theorem 3.1. *Let $\sigma = A_{2g}A_{2g-1}\dots A_2A_1$ be the order $4g+2$ torsion element and $\tau \circ B$ be the order-two element as we describe above. Then for $g \geq 5$ we have $Mod^{\pm}(S_g) =< \sigma, \tau \circ B >$.*

Remark 3.2. *From theorem 3.1 we can also deduce that $Mod^{\pm}(S_g)$ can be generated by 3 symmetries when $g \geq 5$. The reason is that the order $4g+2$ torsion element σ is like the rotation of the $(4g+2)$-gon. In the dihedral group, the order $4g+2$ rotation is the product of two reflections.*

Proof of Theorem 3.1. Denote the subgroup generated by σ and $\tau \circ B$ as G. We proof $G = \mathrm{Mod}^{\pm}(S_g)$ by four steps:

Step 1. For integers i, k satisfying $k \in \{1, 2, \dots, 4g+1\} - \{1, 2, 3, 2g-2, 2g, 2g+2, 2g+4, 4g-1, 4g, 4g+1\}$, b_i is disjoint from b_{i+k} and $B_i\, B_{i+k}^{-1} \in G$.

Step 2. For integers m, n satisfying $m \notin \{n, n+4\}$, a_m is disjoint from b_n and $A_m\, B_n^{-1} \in G$.

Step 3. Using lantern relation, prove that for all k, $A_k \in G$.

Step 4. $G = \mathrm{Mod}^{\pm}(S_g)$.

The proof of Step 1:

For $k \in \{1, 2, \dots, 4g+1\} - \{1, 2, 3, 2g-2, 2g, 2g+2, 2g+4, 4g-1, 4g, 4g+1\}$, $b_0 = b$ is obviously disjoint from b_k (see Figure 2 and Figure 3). B_0 and B_k commute. Consider the element $\sigma^{-k}\,(\tau \circ B)\,\sigma^{-k}\,(\tau \circ B)$. In the dihedral group of the $(4g+2)$-gon, $\sigma^{-k}\,\tau\,\sigma^{-k}\,\tau$ is the identity element. After gluing the $(4g+2)$-gon to a genus g surface and adding the Dehn twist B, we can easily check that $\sigma^{-k}\,(\tau \circ B)\,\sigma^{-k}\,(\tau \circ B) = B_0 B_k^{-1}$. Then conjugating it with σ^{-i}, we have $B_i B_{i+k}^{-1} \in G$.

The proof of Step 2:

By the conjugacy relation, if there exists pair of disjoint curves $(b_i, b_{i+k}$ and $\varphi \in G$ such that $\varphi : (b_i, b_{i+k}) \mapsto (a_m, b_n)$ for some i, k, m, n, then $A_m B_n^{-1}$ is in G.

Suppose first that g is odd. See figure 5.



Figure 5

By step 1, $B_{4g+1}^{-1}B_{2g+2}$ is in G. If $g \geq 5$, b_{2g+2} is disjoint from b_{2g-3}, b_{2g-4}. Both $B_{2g-4}^{-1}B_{2g+2}$ and $B_{2g-3}^{-1}B_{2g+2}$ are in G. $(B_{2g-4}^{-1}B_{2g+2})(B_{2g-3}^{-1}B_{2g+2})$ maps the pair of curves (b_{4g+1}, b_{2g+2}) to the pair of curves (c_1, b_{2g+2}) as in Figure 5 (2). Hence $C_1^{-1}B_{2g+2}$ is in G. Now $B_{2g+2}^{-1}B_{2g-3}$ is also in G. So $C_1^{-1}B_{2g-3}$ is in G. $(B_6^{-1}B_{2g-3})(B_7^{-1}B_{2g-3}) \ldots (B_{2g-8}^{-1}B_{2g-3})(B_{2g-7}^{-1}B_{2g-3})$ maps the pair of curves (c_1, b_{2g-3}) to the pair of curves (c_2, b_{2g-3}) as in Figure 5 (4). Then we have $C_2^{-1}B_{2g-3}$ is in G. $(B_{2g-2}^{-1}B_3)(B_{2g-3}^{-1}B_2)(B_{2g-3}^{-1}B_0)$ maps the pair of curves (c_2, b_{2g-3}) to the pair of curves (a_3, b_{2g-3}). So we get that $A_3^{-1}B_{2g-3}$ is in G. Its inverse $A_3B_{2g-3}^{-1}$ is also in G.

Suppose now that g is even. See figure 6.



Figure 6

In this case, the proof of $C_1^{-1}B_{2g-3} \in G$ is the same as the odd genus case. $(B_4^{-1}B_{2g-3})(B_5^{-1}B_{2g-3}) \ldots (B_{2g-8}^{-1}B_{2g-3})(B_{2g-7}^{-1}B_{2g-3})$ maps the pair of curves (c_1, b_{2g-3}) to the pair of curves (c_2, b_{2g-3}) as in figure 6 (2). $B_{2g-3}^{-1}B_0$ maps the pair of curves (c_2, b_{2g-3}) to the pair of curves (a_4, b_{2g-3}). So we get that $A_4^{-1}B_{2g-3}$ is in G. Its inverse $A_4B_{2g-3}^{-1}$ is also in G.

Again by the conjugacy relation, σ^{2g-3} maps (a_3, b_{2g-3}) to (a_{2g+8}, b) when g is odd and maps (a_4, b_{2g-3}) to (a_{2g+9}, b) when g is even. Now we have $A_i B^{-1} \in G$ for some i. $(\tau \circ B)\, \sigma^{-k}\, (\tau \circ B)(A_i B^{-1}) = A_{i-k} B^{-1}$. Take all possible k and conjugate by σ^n, we have all $A_m B_n^{-1} \in G$ for $m \notin \{n, n+4\}$. The reason we need $m \notin \{n, n+4\}$ is that when $m = n$ or $m = n+4$, the intersection number of a_m and b_n is not zero.

The proof of Step 3:

See figure 7. There is a natural lantern lying on the surface, bounded by a_1, a_3, a_5 and f. By the lantern relation, we have $B_0 B_2 E = A_1 A_3 A_5 F$, or $A_1 = (B_0 A_3^{-1})(B_2 A_5^{-1})(E F^{-1})$, where e and f are the curves showed in figure 7. By the result of step 2, $B_0 A_3^{-1}$ and $B_2 A_5^{-1}$ are in G. We only need to prove EF^{-1} is also in G.



Figure 7

When $g \geq 5$, b_{2g-3} is disjoint from $b_0, \ldots, b_3, a_1, \ldots, a_6, e, f$. $(B_3^{-1} B_{2g-3})(A_6 B_{2g-3}^{-1})(A_5 B_{2g-3}^{-1})(A_4 B_{2g-3}^{-1})$ maps the pair of curves (b_{2g-3}, b_0) to (b_{2g-3}, f). By step 1, $B_{2g-3} B_0^{-1}$ is in G. So $B_{2g-3} F^{-1} \in G$. $(A_2 B_{2g-3}^{-1})(A_1 B_{2g-3}^{-1})(A_4^{-1} B_{2g-3})(B_1 B_{2g-3}^{-1})$ maps the pair of curves (b_{2g-3}, a_5) to (b_{2g-3}, e). By step 2, $B_{2g-3} A_5^{-1}$ is in G. So $B_{2g-3} E^{-1} \in G$ and $EF^{-1} = (E B_{2g-3}^{-1})(B_{2g-3} F^{-1}) \in G$.

The proof of Step 4:

$A_1 \in G$ together with $A_1 B^{-1} \in G$ means that $B \in G$. Now all the A_i's and B_j's are in G. These are Humphries' generators of $\mathrm{Mod}(S_g)$. So G contains $\mathrm{Mod}(S_g)$. Finally, one of the generator $\tau \circ B$ is an orientation-reversing mapping class. Hence $G = \mathrm{Mod}^{\pm}(S_g)$.

□

References

[1] T. E. Brendle and B. Farb. Every mapping class group is generated by 3 torsion elements and by 6 involutions. J. Algebra 278 (2004), 187C198.

[2] J. Brock and D. Margalit. Weil-Petersson isometries via the pants complex. Proc. Amer. Math. Soc., 135(3):795C803 (electronic), 2007.

[3] M. Dehn. Papers on group theory and topology (Springer-Verlag, New York, 1987) (Die Gruppe der Abbildungsklassen, Acta Math. Vol. 69 (1938), 135C206).

[4] B. Farb and D. Marglait. A Primer on Mapping Class Groups. Princeton Math. Ser., (Princeton University Press, 2012), 623C658.

[5] S. P. Humphries. Generators for the mapping class group. Topology of low-dimensional manifolds. Proc. Second Sussex Conf. Chelwood Gate 1977 Lecture Notes in Math. 722 (Springer, 1979), 44C47.
[6] N. V. Ivanov. Automorphism of complexes of curves and of Teichmuller spaces. Internat. Math. Res. Notices, (14):651C666, 1997.
[7] D. Johnson. The structure of Torelli group I: A finite set of generators for I. Ann. of Math. 118 (1983), 423C442.
[8] M. Kassabov. Generating Mapping Class Groups by Involutions. arXiv:math.GT/0311455 v1 25 Nov 2003.
[9] M. Korkmaz. Generating the surface mapping class group by two elements. Trans. Amer. Math. Soc. 357 (2005), 3299C3310.
[10] M. Korkmaz. Minimal generating sets for the mapping class group of a surface. Handbook of Teichmüller theory. Volume III, 441C463, IRMA Lect. Math. Theor. Phys., 17, Eur. Math. Soc., Zürich, 2012.
[11] W. B. R. Lickorish. A finite set of generators for the homeotopy group of a 2-manifold. Proc. Camb. Phils. Soc. 60 (1964), 769C778.
[12] F. Luo. Torsion Elements in the Mapping Class Group of a Surface. arXiv:math.GT/0004048 v1 8 Apr 2000.
[13] D. Margalit. Automorphisms of the pants complex. Duke Math. J., 121(3):457C479, 2004.
[14] N. Monden. Generating the mapping class group by torsion elements of small order. Mathematical Proceedings of the Cambridge Philosophical Society, 154 (2013), pp 41-62, doi:10.1017/S0305004112000357.
[15] J. D. McCarthy and A. Papadopoulos. Involutions in surface mapping class groups.Enseign. Math.(2) 33 (1987), no. 3C4, 275C290. 456.
[16] M. Stukow. The extended mapping class group is generated by 3 symmetries. C. R. Math. Acad. Sci. Paris 338 (2004), no. 5, 403C406.
[17] B. Wajnryb. Mapping class group of a surface is generated by two elements. Topology, 35(2), 377-383, 1996.

South China University of Technology, Guangzhou 510640, P.R.China

E-mail address: scxmdu@scut.edu.cn